

CONTENTS

More detailed tables of contents can be found within the various parts of the compendium.


Acknowledgments	ii
Introduction	iii
Maps of HIV and SIV Genomes	iv
Landmarks of the Genome	v
PART I. REVIEWS	
Tat, a Novel Regulator of HIV Transcription and Latency	2
<i>Jonathan Karn</i>	
An Overview of HIV-1 Co-receptor Function and Its Inhibitors	19
<i>Emmanuel G. Cormier and Tatjana Dragic</i>	
An Overview of the Molecular Phylogeny of Lentiviruses	35
<i>Brian T. Foley</i>	
Sequence Homology Search Tools on the World Wide Web	44
<i>Ian Holmes</i>	
Recombinant HIV Sequences: Their Role in the Global Epidemic	54
<i>Martine Peeters</i>	
Reagents for HIV/SIV Vaccine Studies	73
<i>Rama Thakallapally and Carla Kuiken</i>	
Clinical Trials of HIV Vaccines	82
<i>Barney S. Graham</i>	
Mutations in Retroviral Genes Associated with Drug Resistance	106
<i>Urvi Parikh, Jennifer Hammond, Charles Calef, Brendan Larder,</i> <i>Raymond Schinazi, John W. Mellors</i>	
PART II. HIV-1/SIVcpz NUCLEOTIDE ALIGNMENTS	
Contents	163
Introduction	164
Table of HIV-1/SIVcpz Sequences in the Nucleotide Alignment	167
Notes on full-length HIV-1/SIVcpz Sequences in the Nucleotide Alignment	170
Nucleotide Alignment of HIV-1/SIVcpz Complete Genomes	188
PART III. HIV-1/HIV-2/SIV NUCLEOTIDE ALIGNMENTS	
Contents	355
Introduction	355
Table of Primate Lentiviruses Sequences in the Nucleotide Alignment	357
Nucleotide Alignment of Primate Lentiviral Complete Genomes	358
PART IV. HIV-1/SIVcpz AMINO ACID ALIGNMENTS	
Contents	457
Introduction	
Table of HIV-1/SIVcpz sequences in the Amino Acid Alignments	460
Amino Acid Alignments of HIV-1/SIVcpz	470
PART V. HIV-2/SIV AMINO ACID ALIGNMENTS	
Contents	531
Introduction	532
Table of HIV-2/SIV sequences in the Amino Acid Alignments	533
Amino Acid Alignments of HIV-2/SIV	537
PART VI. Other SIV AMINO ACID ALIGNMENTS	
Contents	567
Table of other SIV sequences in the Amino Acid Alignments	568
Amino Acid Alignments of Other SIV	570

ACKNOWLEDGMENTS

The HIV Sequence Database and Analysis Project is funded by the Vaccine and Prevention Research Program of the AIDS Division of the National Institute of Allergy and Infectious Diseases (Dr. James Bradac, Project Officer) through an interagency agreement with the U.S. Department of Energy.

We thank our editors, the many researchers who have made their sequences available prior to publication, and authors who help by contributing to our review section.

The Cover



A schematic representation of the activation mechanism of latent proviruses by NF- κ B and Tat during T-cell activation. From: Karn J, Tat, a novel regulator of HIV transcription and latency, Page 2 of this Compendium.

Citing this publication

We have simplified the name of this, our annual publication. Formerly known as “Human Retroviruses and AIDS” it should now be cited simply as *HIV Sequence Compendium 2000*, Kuiken C, Foley B, Hahn B, Marx P, McCutchan F, Mellors J, Mullins J, Wolinsky S, and Korber B, editors. Published by Theoretical Biology and Biophysics Group, Los Alamos National Laboratory.

Introduction

This Compendium is an annual printed summary of the data contained in the HIV sequence database. In these compendia we try to present a judicious selection of the data in such a way that it is of maximum utility to HIV researchers. Traditionally, we present the sequence data themselves in the form of alignments: a comprehensive alignment of a selection of all full-length genomes the database contains (a lot of LAI-like sequences, for example, have been omitted because they are so similar that they bias the alignment) of HIV-1/SIVcpz (Section I) and a combined HIV-1/HIV-2/SIV whole genome alignment (Section II); amino acid alignments for HIV-1/SIV-cpz, HIV-2/SIV, and SIVagm. The HIV-2/SIV and SIVagm amino acid alignments are separate because the genetic distances between these groups are so great that presenting them in one alignment would make them very elongated because of the large number of gaps that have to be inserted. As always, tables with extensive background information gathered from the literature accompany the whole genome alignments.

The collection of whole-gene sequences in the database is now large enough that we have abundant representation of most subtypes (excluding H and J). For most other subtypes, and especially for subtype B, a large number of sequences that span entire genes were not included in the printed alignments to conserve space. A more complete version of all alignments is available on our website, <http://hiv-web.lanl.gov> Importantly, all these alignments have been edited to include only one sequence per person, based on phylogenetic trees that were created for all of them, as well as the literature. At the request of many users, we have re-inserted the consensus sequences for each subtype, unless there are fewer than five sequences representing it. In the alignments we have also included the ‘Circulating Recombinant Forms’, mosaic genomes that have epidemiological significance (see the nomenclature chapter for more on CRFs). Finally, for all amino acid alignments we have decided to combine the annotation tables into one, because of the increasing redundancy in the separate tables. In addition to sequence information (accession numbers, references) the new table lists which regions of the sequence are represented in the alignments.

We have made an effort to bring the HIV-2/SIV and SIVagm alignments up-to-date as well. We have created an entirely new HIV-1/HIV-2/SIV alignment that is much improved over the previous version; it can be accessed via our website. Because of the frequency of redundant information, we have decided to merge the gene tables into one large table for each alignment section; we hope you will find these tables easy to use.

In the Reviews section, along with the previously mentioned chapter on HIV nomenclature, you will find a very clearly written and concise overview of the functions of TAT; a review of lentivirus phylogeny and evolution; an overview of recombinants and their role in the epidemic; a concise and lucid chapter on protein search tools on the internet; an overview of HIV-1 coreceptors and coreceptor inhibitors; and a very thorough review on progress in SIV and HIV vaccines. In addition, we present updated versions of the customary reviews of coreceptor usage, drug resistance, and SIV/SHIV vaccine reagents. Reprints of all reviews are available from our website in the form of both HTML and PDF files.

As always, we are open to complaints and suggestions for improvement. With the effort that goes into producing these volumes, we sincerely hope they will be widely used by the research community. Inquiries and comments regarding the Compendium should be addressed to:

Dr. Carla Kuiken

Theoretical Division, T-10, MS K710, LANL, Los Alamos, NM 87545

Ph: (505)-665—6463; fax: (505)-665—3493; e-mail: kuiken@t10.lanl.gov

HIV/SIV PROTEINS			
NAME	SIZE	FUNCTION	LOCALIZATION
Gag MA	p17	membrane anchoring; env interaction; nuclear transport of viral core. (myristylated protein)	virion
CA	p24	core capsid	virion
NC	p7	nucleocapsid, binds RNA	virion
	p6	binds Vpr	virion
Protease (PR)	p15	gag/pol cleavage and maturation	virion
Reverse transcriptase (RT), RNase H	p66 p51	reverse transcription, RNase H activity	virion
Integrase (IN)		DNA provirus integration	virion
Env	gp120 gp41	external viral glycoproteins bind to CD4 and secondary receptors	plasma membrane, virion envelope
Tat	p16/p14	viral transcriptional transactivator	primarily in nucleolus/nucleus
Rev	p19	RNA transport, stability and utilization factor (phosphoprotein)	primarily in nucleolus/nucleus shuttling between nucleolus and cytoplasm
Vif	p23	promotes virion maturation and infectivity	cytoplasm (cytosol, membranes) virion
Vpr	p10-15	promotes nuclear localization of preintegration complex, inhibits cell division, arrests infected cells at G2/M	virion, nucleus (nuclear membrane?)
Vpu	p16	promotes extracellular release of viral particles; degrades CD4 in the ER; (phosphoprotein only in HIV-1 and SIVcpz)	integral membrane protein
Nef	p27-p25	CD4 and class I downregulation (myristylated protein)	plasma membrane, cytoplasm (virion?)
Vpx	p12-16	vpr homolog (not in HIV-1, only in HIV-2 and SIV)	virion (nucleus?)

LANDMARKS:

HIV GENOMIC STRUCTURAL ELEMENTS

- LTR** Long terminal repeat, the DNA sequence flanking the genome of integrated proviruses. It contains important regulatory regions, especially those for transcription initiation and polyadenylation.
- TAR** Target sequence for viral transactivation, the binding site for Tat protein and for cellular proteins; consists of approximately the first 45 nucleotides of the viral mRNAs in HIV-1 (or the first 100 nucleotides in HIV-2 and SIV.) TAR RNA forms a hairpin stem-loop structure with a side bulge; the bulge is necessary for Tat binding and function.
- RRE** Rev responsive element, an RNA element encoded within the env region of HIV-1. It consists of approximately 200 nucleotides (positions 7327 to 7530 from the start of transcription in HIV-1, spanning the border of gp120 and gp41). The RRE is necessary for Rev function; it contains a high affinity site for Rev; in all, approximately seven binding sites for Rev exist within the RRE RNA. Other lentiviruses (HIV-2, SIV, visna, CAEV) have similar RRE elements in similar locations within env, while HTLVs have an analogous RNA element (RXRE) serving the same purpose within their LTR; RRE is the binding site for Rev protein, while RXRE is the binding site for Rex protein. RRE (and RXRE) form complex secondary structures, necessary for specific protein binding.
- CRS** Cis-acting repressive sequences postulated to inhibit structural protein expression in the absence of Rev. One such site was mapped within the pol region of HIV-1. The exact function has not been defined; splice sites have been postulated to act as CRS sequences.
- INS** Inhibitory/Instability RNA sequences found within the structural genes of HIV-1 and of other complex retroviruses. Multiple INS elements exist within the genome and can act independently; one of the best characterized elements spans nucleotides 414 to 631 in the gag region of HIV-1. The INS elements have been defined by functional assays as elements that inhibit expression posttranscriptionally. Mutation of the RNA elements was shown to lead to INS inactivation and up regulation of gene expression.

GENES AND GENE PRODUCTS

- GAG** The genomic region encoding the capsid proteins (group specific antigens). The precursor is the p55 myristylated protein, which is processed to p17 (MA_{matrix}), p24 (CA_{capsid}), p7 (NucleoCapsid), and p6 proteins, by the viral protease. Gag associates with the plasma membrane where the virus assembly takes place. The 55 kDa Gag precursor is called assemblin to indicate its role in viral assembly.
- POL** The genomic region encoding the viral enzymes protease, reverse transcriptase and integrase. These enzymes are produced as a Gag-pol precursor polyprotein, which is processed by the viral protease; the Gag-pol precursor is produced by ribosome frameshifting at the C-terminus of gag.
- ENV** Viral glycoproteins produced as a precursor (gp160) which is processed to give a noncovalent complex of the external glycoprotein gp120 and the transmembrane glycoprotein gp41. The mature gp120-gp41 proteins are bound by non-covalent interactions and are associated as a trimer on the cell surface. A substantial amount of gp120 can be found released in the medium. gp120 contains the binding site for the CD4 receptor, and the seven transmembrane domain chemokine receptors that serve as co-receptors for HIV-1.
- TAT** Transactivator of HIV gene expression. One of two essential viral regulatory factors (Tat and Rev) for HIV gene expression. Two forms are known, Tat-1 exon (minor form) of 72 amino acids and Tat-2exon (major form) of 86 amino acids. Low levels of both proteins are found in persistently infected cells. Tat has been localized primarily in the nucleolus/nucleus by immunofluorescence. It acts by binding to the TAR RNA element and activating transcription

initiation and/or elongation from the LTR promoter. It is the first eukaryotic transcription factor known to interact with RNA rather than DNA and may have similarities with prokaryotic anti-termination factors. Extracellular Tat can be found and can be taken up by cells in culture.

- REV** The second necessary regulatory factor for HIV expression. A 19 kD phosphoprotein, localized primarily in the nucleolus/nucleus, Rev acts by binding to RRE and promoting the nuclear export, stabilization and utilization of the viral mRNAs containing RRE. Rev is considered the most functionally conserved regulatory protein of lentiviruses. Rev cycles rapidly between the nucleus and the cytoplasm.
- VIF** Viral infectivity factor, a basic protein of typically 23 kD. Promotes the infectivity but not the production of viral particles. In the absence of Vif the produced viral particles are defective, while the cell-to-cell transmission of virus is not affected significantly. Found in almost all lentiviruses, Vif is a cytoplasmic protein, existing in both a soluble cytosolic form and a membrane-associated form. The latter form of Vif is a peripheral membrane protein that is tightly associated with the cytoplasmic side of cellular membranes. Some recent observations suggest that Vif functions late in replication to modulate assembly, budding, and/or maturation the N-terminal half of Vif (N'-Vif) specifically interacts with viral protease.
- VPR** Vpr (viral protein R) is a 96-amino acid (14 kd) protein, which is incorporated into the virion. It interacts with the p6 gag part of the Pr55 gag precursor. Vpr detected in the cell is localized to the nucleus. Proposed functions for Vpr include the targeting the nuclear import of preintegration complexes, cell growth arrest, transactivation of cellular genes, and induction of cellular differentiation. It is found in HIV-1, HIV-2, SIVmac and SIVmnd. It is homologous to the vpx protein.
- VPU** Vpu (viral protein U) is unique to HIV-1 and SIVcpz, a close relative of HIV-1. There is no similar gene in HIV-2 or other SIVs. Vpu is a 16-kd (81-amino acid) type I integral membrane protein with at least two different biological functions: (a) degradation of CD4 in the endoplasmic reticulum, and (b) enhancement of virion release from the plasma membrane of HIV-1-infected cells. Env and Vpu are expressed from a bicistronic mRNA. Vpu probably possesses an N-terminal hydrophobic membrane anchor and a hydrophilic moiety. It is phosphorylated by casein kinase II at positions Ser52 and Ser56. Vpu is involved in env maturation and is not found in the virion. Vpu has been found to increase susceptibility of HIV-1 infected cells to Fas killing.
- NEF** A multifunctional 27-kd myristylated protein produced by an ORF located at the 3' end of the primate lentiviruses. Other forms of Nef are known, including nonmyristylated variants. Nef is predominantly cytoplasmic and associated with the plasma membrane via the myristyl residue linked to the conserved second amino acid (Gly). Nef has also been identified in the nucleus and found associated with the cytoskeleton in some experiments. One of the first HIV proteins to be produced in infected cells, it is the most immunogenic of the accessory proteins. The nef genes of HIV and SIV are dispensable *in vitro*, but are essential for efficient viral spread and disease progression *in vivo*. Nef is necessary for the maintenance of high virus loads and for the development of AIDS in macaques, and viruses with defective Nef have been detected in some HIV-1 infected long term survivors. Nef downregulates CD4, the primary viral receptor, and MHC class I molecules, and these functions map to different parts of the protein. Nef interacts with components of host cell signal transduction and clathrin-dependent protein sorting pathways. It increases viral infectivity. Nef contains PxxP motifs that bind to SH3 domains of a subset of Src kinases and are required for the enhanced growth of HIV but not for the downregulation of CD4.
- VPX** A virion protein of 12 kD found only in HIV-2/SIVmac/SIVsm and not in HIV-1 or SIVagm. This accessory gene is a homolog of HIV-1 vpr, and HIV-2/SIV carry both vpr and vpx. Vpx function in relation to vpr is not fully elucidated; both are incorporated into virions at levels comparable to gag proteins through interactions with Gag p6. Vpx is necessary for efficient replication of SIV in PBMCs. Progression to AIDS and death in SIV-infected animals can occur in the absence of Vpr or Vpx. Double mutant virus lacking both vpr and vpx was attenuated,

whereas the single mutants were not, suggesting a redundancy in the function of Vpr and Vpx related to virus pathogenicity.

STRUCTURAL PROTEINS/VIRAL ENZYMES The products of gag, pol and env genes, which are essential components of the retroviral particle.

REGULATORY PROTEINS Tat and Rev proteins of HIV/SIV and Tax and Rex proteins of HTLVs. They modulate transcriptional and posttranscriptional steps of virus gene expression and are essential for virus propagation.

ACCESSORY OR AUXILIARY PROTEINS Additional virion and non-virion- associated proteins produced by HIV/SIV retroviruses: Vif, Vpr, Vpu, Vpx, Nef. Although the accessory proteins are in general not necessary for viral propagation in tissue culture, they have been conserved in the different isolates; this conservation and experimental observations suggest that their role *in vivo* is very important. Their functional importance continues to be elucidated.

COMPLEX RETROVIRUSES Retroviruses regulating their expression via viral factors and expressing additional proteins (regulatory and accessory) essential for their life cycle.